

On equations defining the Ricci flows of manifolds

Valerii Dryuma*

Institute of Mathematics and Informatics, AS RM,

5 Academiei Street, 2028 Kishinev, Moldova,

e-mail: valery@dryuma.com; valdryum@gmail.com

Abstract

The examples of the Ricci flows of the four-dimensional manifolds which are determined by help of nonlinear differential equations of the type of Monge-Ampere are constructed. Their particular solutions are derived and their properties are discussed.

1 $4D$ model of the Ricci-flow

We study the system of equations

$$\frac{\partial}{\partial t} g_{ij}(\vec{x}, t) = -2R_{ij}(g) \quad (1)$$

describing the Ricci flows of the four dimensional manifolds which are endowed by the metrics of the form

$$ds^2 = A(x, y, t) du^2 + 2 B(x, y, t) du dv + du dx + C(x, y, t) dv^2 + dv dy, \quad (2)$$

where the components of metrics are dependent from two coordinates (x, y) and from the parameter t .

In this case the Ricci-tensor of the metric (1) has a five components

$$R_{uu}, R_{uv}, R_{vv}, R_{ux}, R_{vx}.$$

From the conditions of compatibility of the equations (1) we find that they are reduced to the equation

$$4 \left(\frac{\partial^2}{\partial x^2} h(x, y, t) \right) \frac{\partial^2}{\partial y^2} h(x, y, t) - 4 \left(\frac{\partial^2}{\partial x \partial y} h(x, y, t) \right)^2 - \frac{\partial}{\partial t} h(x, y, t) = 0 \quad (3)$$

and components of the metric (2) take the form

$$B(x, y, t) = \frac{\partial^2}{\partial x \partial y} h(x, y, t), \quad C(x, y, t) = -\frac{\partial^2}{\partial x^2} h(x, y, t), \quad A(x, y, t) = -\frac{\partial^2}{\partial y^2} h(x, y, t). \quad (4)$$

*Work supported in part by Grant RFFI, Russia-Moldova

Thus the study of the properties of the Ricci flow on a 4-dimensional manifold with the metric (2) with the coefficients (4) lead to integration of the equation (3). Nyis equation is the first order relatively the parameter τ and has form of the Monge-Ampere equation relatively of the variables x, y

2 Particular solutions

Let us consider some elementary solutions of equation (3)

After the substitution of the form

$$h(x, y, t) = H \left(\frac{x}{y} - t \right) y^4$$

the equation (3) takes the form

$$48 \left(D^{(2)} \right) (H)(\eta) H(\eta) - 36 (D(H)(\eta))^2 + D(H)(\eta) = 0, \quad (5)$$

where

$$\eta = \frac{x}{y} - t.$$

Solution of the equation (4) may be present in form

$$\eta(H) = -C2 - 144 \frac{\sqrt[4]{H}}{K^3} + \left(-48 \ln(1 - \sqrt[4]{H}K) + 24 \ln(1 + \sqrt[4]{H}K + \sqrt{H}K^2) + 48 \sqrt{3} \arctan\left(\frac{\sqrt{3}\sqrt[4]{H}K}{2 + \sqrt[4]{H}K}\right) \right) K^{-4},$$

where K and $-C2$ are constants.

Remark that if the value $K = -1$ the function $\eta(H)$ has a break.

The simplest singular solution of (4) is

$$h(x, y, t) = \frac{\left(6 \frac{B^2 x^4}{k} + -C1 \sqrt{x} \cos(1/2 \sqrt{23} \ln(x)) + -C2 \sqrt{x} \sin(1/2 \sqrt{23} \ln(x)) + y B x^3 + 1/24 y^2 k x^2 \right)}{(2 k t + -C4)^{-1}}.$$

More complicated solution has the form

$$h(x, y, t) = -1/2 \frac{-C1^2 \mu \sqrt{-C1 - c_1} \sqrt{2} \tan(1/2 \sqrt{-C1 - c_1} (x - t) \sqrt{2})}{-c_1^2} + \frac{\mu -C1 y}{-c_1} -$$

$$-1/4 \frac{\mu \sqrt{-C1 - c_1} \sqrt{2} y^2}{-c_1 \tan(1/2 \sqrt{-C1 - c_1} (x - t) \sqrt{2}) -C1}.$$

A more general solutions of the equation (4) are derived by the method of parametric representations of functions and their derivatives [1-2].

Let us apply its to the equation (4).

After the change of the variables, the function and its derivative

$$h(x, y, t) \rightarrow u(x, \tau, t), \quad y \rightarrow v(x, \tau, t), \quad h_x \rightarrow u_x - \frac{u_\tau}{v_\tau} v_x = p, \quad h_t \rightarrow u_t - \frac{u_\tau}{v_\tau} v_t = q,$$

$$h_y \rightarrow \frac{u_\tau}{v_\tau} = r, \quad h_{xx} \rightarrow p_x - \frac{p_\tau}{v_\tau} v_x, \quad h_{xy} \rightarrow \frac{p_\tau}{v_\tau} = q_x - \frac{q_\tau}{v_\tau} v_x, \quad h_{yy} \rightarrow \frac{r_\tau}{v_\tau} \quad (6)$$

we derive from the equation

$$\left(\frac{\partial^2}{\partial x^2} h(x, y, t) \right) \frac{\partial^2}{\partial y^2} h(x, y, t) - \left(\frac{\partial^2}{\partial x \partial y} h(x, y, t) \right)^2 - \frac{\partial}{\partial t} h(x, y, t) = 0 \quad (7)$$

the relation between the functions $u(x, \tau, t)$ and $v(x, \tau, t)$, where variable τ is considered as parameter

$$\begin{aligned} & - \left(\frac{\partial^2}{\partial x^2} u(x, \tau, t) \right) \left(\frac{\partial}{\partial \tau} v(x, \tau, t) \right) \left(\frac{\partial}{\partial \tau} u(x, \tau, t) \right) \frac{\partial^2}{\partial \tau^2} v(x, \tau, t) - \\ & - \left(\frac{\partial}{\partial \tau} u(x, \tau, t) \right) \left(\frac{\partial^2}{\partial x^2} v(x, \tau, t) \right) \left(\frac{\partial}{\partial \tau} v(x, \tau, t) \right) \frac{\partial^2}{\partial \tau^2} u(x, \tau, t) + \\ & + \left(\frac{\partial}{\partial \tau} u(x, \tau, t) \right)^2 \left(\frac{\partial^2}{\partial x^2} v(x, \tau, t) \right) \frac{\partial^2}{\partial \tau^2} v(x, \tau, t) - \\ & - \left(\frac{\partial^2}{\partial \tau \partial x} u(x, \tau, t) \right)^2 \left(\frac{\partial}{\partial \tau} v(x, \tau, t) \right)^2 + \left(\frac{\partial^2}{\partial x^2} u(x, \tau, t) \right) \left(\frac{\partial}{\partial \tau} v(x, \tau, t) \right)^2 \frac{\partial^2}{\partial \tau^2} u(x, \tau, t) + \\ & + 2 \left(\frac{\partial^2}{\partial \tau \partial x} u(x, \tau, t) \right) \left(\frac{\partial}{\partial \tau} v(x, \tau, t) \right) \left(\frac{\partial}{\partial \tau} u(x, \tau, t) \right) \frac{\partial^2}{\partial \tau \partial x} v(x, \tau, t) - \\ & - \left(\frac{\partial}{\partial \tau} u(x, \tau, t) \right)^2 \left(\frac{\partial^2}{\partial \tau \partial x} v(x, \tau, t) \right)^2 - \left(\frac{\partial}{\partial t} u(x, \tau, t) \right) \left(\frac{\partial}{\partial \tau} v(x, \tau, t) \right)^4 + \\ & + \left(\frac{\partial}{\partial \tau} u(x, \tau, t) \right) \left(\frac{\partial}{\partial t} v(x, \tau, t) \right) \left(\frac{\partial}{\partial \tau} v(x, \tau, t) \right)^3 = 0. \end{aligned} \quad (8)$$

Note that the relation (8) under the condition $v(x, \tau, t) = \tau$ is equivalent to the equation

$$\frac{\partial^2}{\partial x^2} u(x, t, \tau) \frac{\partial^2}{\partial t^2} u(x, t, \tau) - \left(\frac{\partial^2}{\partial t \partial x} u(x, t, \tau) \right)^2 - \frac{\partial}{\partial \tau} u(x, t, \tau) = 0$$

similar to equation (7) and transformed into an equation of more general form if the functions $u(x, \tau, t)$, $(v(x, \tau, t))$ are dependent.

For example under the condition

$$u(x, \tau, t) = \tau \frac{\partial}{\partial \tau} \omega(x, \tau, t) - \omega(x, \tau, t), \quad v(x, \tau, t) = \frac{\partial}{\partial \tau} \omega(x, \tau, t)$$

from the (8) we obtain the p.d.e. for the function $\omega(x, \tau, t)$

$$-\frac{\partial^2}{\partial x^2} \omega(x, \tau, t) + \left(\frac{\partial^2}{\partial \tau^2} \omega(x, \tau, t) \right) \frac{\partial}{\partial t} \omega(x, \tau, t) = 0. \quad (9)$$

The other condition

$$v(x, \tau, t) = \tau \frac{\partial}{\partial \tau} \omega(x, \tau, t) - \omega(x, \tau, t), \quad u(x, \tau, t) = \frac{\partial}{\partial \tau} \omega(x, \tau, t)$$

lead to the equation

$$\frac{\partial^2}{\partial x^2} \omega(x, \tau, t) + \left(\frac{\partial^2}{\partial \tau^2} \omega(x, \tau, t) \right) \tau^3 \frac{\partial}{\partial t} \omega(x, \tau, t) = 0. \quad (10)$$

Let us consider some solutions of the equation (9).

Under the substitution

$$\omega(x, \tau, t) = A(x - kt, \tau)$$

it is reduced at the equation

$$\frac{\partial^2}{\partial \eta^2} A(\eta, \tau) + k \left(\frac{\partial^2}{\partial \tau^2} A(\eta, \tau) \right) \frac{\partial}{\partial \eta} A(\eta, \tau) = 0 \quad (11)$$

where $\eta = x - kt$.

This equation is reduced after the Legendre transformation to the Euler-Trikomi equation and therefore its solutions depend radically on the sign of the parameter k .

Each solution (11) corresponds to the solution of the original equation (8) which is obtained by eliminating the parameter τ from the correlations

$$y - \tau \frac{\partial}{\partial \tau} \omega(x, \tau, t) + \omega(x, \tau, t) = 0, \quad h(x, y, t) - \frac{\partial}{\partial \tau} \omega(x, \tau, t) = 0. \quad (12)$$

Here is an example of.

3 Finite-dimensional reduction

After the substitutions

$$A(x, y, t) = A_0(t) + A_1(t)x + A_2(t)y + A_3(t)x^2 + A_4(t)yx + A_5(t)y^2,$$

$$B(x, y, t) = B_0(t) + B_1(t)x + B_2(t)y + B_3(t)x^2 + B_4(t)xy + B_5(t)y^2,$$

$$C(x, y, t) = C_0(t) + C_1(t)x + C_2(t)y + C_3(t)x^2 + C_4(t)xy + C_5(t)y^2,$$

$$B_4(t) = -2 A_3(t), \quad A_4(t) = -2 B_5(t), \quad C_4(t) = -2 B_3(t), \quad C_5(t) = A_3(t)$$

the metric (2) takes the form

$$\begin{aligned} ds^2 = & \left(A_0(t) + A_1(t)x + A_2(t)y + A_3(t)x^2 - 2 B_5(t)yx + A_5(t)y^2 \right) du^2 + \\ & + 2 \left(B_0(t) + B_1(t)x + B_2(t)y + B_3(t)x^2 - 2 A_3(t)xy + B_5(t)y^2 \right) du dv + \\ & + \left(C_0(t) + C_1(t)x + C_2(t)y + C_3(t)x^2 - 2 B_3(t)xy + A_3(t)y^2 \right) dv^2 + dx du + dy dv \end{aligned} \quad (13)$$

and from the system of equations (1) is obtained system of ODE's

$$\frac{d}{dt} B_3(t) = 24 B_3(t) A_3(t) - 24 C_3(t) B_5(t),$$

$$\frac{d}{dt} C_3(t) = -48 C_3(t) A_3(t) + 48 (B_3(t))^2,$$

$$\begin{aligned}
\frac{d}{dt}A_3(t) &= -8 C_3(t)A_5(t) + 24 (A_3(t))^2 - 16 B_3(t)B_5(t), \\
\frac{d}{dt}B_5(t) &= 24 A_3(t)B_5(t) - 24 B_3(t)A_5(t), \\
\frac{d}{dt}A_5(t) &= -48 A_5(t)A_3(t) + 48 (B_5(t))^2
\end{aligned} \tag{14}$$

on the functions $A_3(t)$, $B_3(t)$, $C_3(t)$, $A_5(t)$, $B_5(t)$ from which expressed the rest coefficients of the metric (13).

The system of equations (14) has a first integral M of the type

$$C_3(t)B_5(t)^2 + M C_3(t)^2 B_5(t) - A_5(t)B_3(t)^2 + 2 M B_3(t)^3 - 3 M C_3(t)A_3(t)B_3(t) = 0$$

with the help of which the order can be reduced.

In result we get the system with parameter

$$\begin{aligned}
\frac{d}{dt}B_3(t) &= 24 B_3(t)A_3(t) - 24 C_3(t)B_5(t), \quad \frac{d}{dt}C_3(t) = -48 C_3(t)A_3(t) + 48 (B_3(t))^2, \\
B_3(t)^2 \frac{d}{dt}A_3(t) &= -8 C_3(t)^2 B_5(t)^2 - 8 M C_3(t)^3 B_5(t) - 16 C_3(t)M B_3(t)^3 + \\
&\quad + 24 M C_3(t)^2 A_3(t)B_3(t) + 24 A_3(t)^2 B_3(t)^2 - 16 B_5(t)B_3(t)^3, \\
B_3(t) \frac{d}{dt}B_5(t) &= 24 A_3(t)B_5(t)B_3(t) - 24 C_3(t)B_5(t)^2 - 24 M C_3(t)^2 B_5(t) - \\
&\quad - 48 M B_3(t)^3 + 72 M C_3(t)A_3(t)B_3(t).
\end{aligned}$$

4 Monge-Ampere flows

Theorem 1 *The flows of Ricci of the 4D- manifolds which are endowed by the metric with local coordinates (x, y, z, t) the components of which are dependent from two coordinates (x, t) and from the parameter τ*

$$\begin{aligned}
ds^2 &= \left(\frac{\partial^2}{\partial x^2} f(x, t, \tau) \right) dx^2 + 2 \left(\frac{\partial^2}{\partial t \partial x} f(x, t, \tau) \right) dx dt + \left(\frac{\partial^2}{\partial t^2} f(x, t, \tau) \right) dy^2 + \\
&\quad + 2 \left(\frac{\partial^2}{\partial t \partial x} f(x, t, \tau) \right) dy dz + \left(\frac{\partial^2}{\partial x^2} f(x, t, \tau) \right) dz^2 + \left(\frac{\partial^2}{\partial t^2} f(x, t, \tau) \right) dt^2
\end{aligned}$$

is defined by the equation

$$\frac{\partial}{\partial \tau} f(x, t, \tau) = \ln \left(\frac{\partial^2}{\partial t^2} f(x, t, \tau) \frac{\partial^2}{\partial x^2} f(x, t, \tau) - \left(\frac{\partial^2}{\partial t \partial x} f(x, t, \tau) \right)^2 \right) \tag{15}$$

Let us consider some solutions of the equation (15).

With aim of convenience we rewrite if in the form

$$\left(\frac{\partial^2}{\partial x^2} f(x, y, \tau) \right) \frac{\partial^2}{\partial y^2} f(x, y, \tau) - \left(\frac{\partial^2}{\partial x \partial y} f(x, y, \tau) \right)^2 - e^{\frac{\partial}{\partial \tau} f(x, y, \tau)} = 0, \tag{16}$$

where the variable t is changed on y .

After the (u, v) -transformation with condition $u(x, t, \tau) = t$ the equation (16) takes the form

$$\left(\frac{\partial^2}{\partial x^2}v(x, t, \tau)\right)\frac{\partial^2}{\partial t^2}v(x, t, \tau) - \left(\frac{\partial^2}{\partial t\partial x}v(x, t, \tau)\right)^2 - e^{-\frac{\partial}{\partial t}v(x, t, \tau)}\left(\frac{\partial}{\partial t}v(x, t, \tau)\right)^4 = 0.$$

Particular solutions of this equation are obtained with the help of additional conditions. For example in the case

$$\frac{\partial}{\partial \tau}v(x, t, \tau) = \frac{\partial}{\partial t}v(x, t, \tau)$$

we find that the function $v(x, t, \tau)$ has the form

$$v(x, t, \tau) = {}_2F1(x, \tau + t) = h(x, \eta),$$

where the function $h(x, \eta)$ satisfies the equation

$$\left(\frac{\partial^2}{\partial \eta^2}h(x, \eta)\right)\frac{\partial^2}{\partial x^2}h(x, \eta) - \left(\frac{\partial^2}{\partial \eta\partial x}h(x, \eta)\right)^2 - e^{-1}\left(\frac{\partial}{\partial \eta}h(x, \eta)\right)^4 = 0. \quad (17)$$

After the (u, v) -transformation with the conditions

$$v(x, t) = t\frac{\partial}{\partial t}\omega(x, t) - \omega(x, t), \quad u(x, t) = \frac{\partial}{\partial t}\omega(x, t)$$

the equation (17) is reduced to the linear equation

$$\frac{\partial^2}{\partial x^2}\omega(x, t) + \left(\frac{\partial^2}{\partial t^2}\omega(x, t)\right)e^{-1} = 0$$

with general solution

$$\omega(x, t) = {}_2F1(t - i\sqrt{e^{-1}}x) + {}_2F2(t + i\sqrt{e^{-1}}x)$$

containing two arbitrary functions.

With the help of the function $\omega(x, t)$ we can obtain large class of solutions of the equation (16) in parametric form.

For example, in the case

$${}_2F1(t - i\sqrt{e^{-1}}x) = \cosh(+ - i\sqrt{e^{-1}}x){}_2F2(t + i\sqrt{e^{-1}}x) = \sinh(t + i\sqrt{e^{-1}}x)$$

we get

$$\omega(x, t) = \cosh(t) \cos(\sqrt{e^{-1}}x) + \sinh(t) \cos(\sqrt{e^{-1}}x)$$

and elimination of the parameter t from the relations

$$\eta - t \cos(e^{-1/2}x) \cosh(t) - t \sinh(t) \cos(e^{-1/2}x) + \cosh(t) \cos(e^{-1/2}x) + \sinh(t) \cos(e^{-1/2}x) = 0,$$

$$h(x, \eta) - \cosh(t) \cos(e^{-1/2}x) - \sinh(t) \cos(e^{-1/2}x)$$

lead to the function

$$h(x, \eta) = e^{\text{Lambert } W\left(\frac{\eta e^{-1}}{\cos(e^{-1/2}x)}\right)+1} \cos(e^{-1/2}x)$$

which is solution of the equation (17).

With the help of solutions of the equation (17) can be constructed solutions of the equation (16).

5 More general solution

The equation (16) after the (u, v) -transformation with conditions

$$-\frac{\left(\frac{\partial}{\partial \tau} u(x, t, \tau)\right) \frac{\partial}{\partial t} v(x, t, \tau) + \left(\frac{\partial}{\partial t} u(x, t, \tau)\right) \frac{\partial}{\partial \tau} v(x, t, \tau)}{\frac{\partial}{\partial t} v(x, t, \tau)} = Ax,$$

$$v(x, t, \tau) = -F1(x, u(x, t, \tau) - Ax\tau)$$

takes the form

$$\left(\frac{\partial^2}{\partial \eta^2} h(x, \eta)\right) \frac{\partial^2}{\partial x^2} h(x, \eta) - \left(\frac{\partial^2}{\partial \eta \partial x} h(x, \eta)\right)^2 - e^{Ax} \left(\frac{\partial}{\partial \eta} h(x, \eta)\right)^4 = 0, \quad (18)$$

where

$$\eta = u(x, t, \tau) - Ax\tau,$$

$$h(x, \eta) = -F1(x, u(x, t, \tau) - Ax\tau).$$

It is reduced to the equation

$$e^{Ax} \frac{\partial^2}{\partial t^2} \omega(x, t) + \frac{\partial^2}{\partial x^2} \omega(x, t) = 0 \quad (19)$$

after the (u, v) -transformation with conditions

$$v(x, t) = t \frac{\partial}{\partial t} \omega(x, t) - \omega(x, t), \quad u(x, t) = \frac{\partial}{\partial t} \omega(x, t).$$

Simplest solution of the equation (19) has the form

$$\omega(x, t) = (-C1 \cos(\sqrt{-c_1}t) + -C2 \sin(\sqrt{-c_1}t)) \cdot$$

$$\cdot \left(-C3 \text{Bessel}J(0, 2 \sqrt{-\frac{c_1}{A^2}} \sqrt{e^{Ax}}) + -C4 \text{Bessel}Y(0, 2 \sqrt{-\frac{c_1}{A^2}} \sqrt{e^{Ax}}) \right)$$

and elimination of the parameter t from the relations

$$\eta + t \sin(\sqrt{-c_1}t) \sqrt{-c_1} \text{Bessel}J(0, 2 \sqrt{-c_1} \sqrt{e^x}) + \cos(\sqrt{-c_1}t) \text{Bessel}J(0, 2 \sqrt{-c_1} \sqrt{e^x}) = 0,$$

$$h(x, \eta) + \sin(\sqrt{-c_1}t) \sqrt{-c_1} \text{Bessel}J(0, 2 \sqrt{-c_1} \sqrt{e^x}) = 0$$

gives solution of the equation (18) defined from the relation

$$\eta \sqrt{-c_1} + \arcsin \left(\frac{h(x, \eta)}{\sqrt{-c_1} \text{Bessel}J(0, 2 \sqrt{-c_1} \sqrt{e^x})} \right) h(x, \eta) +$$

$$+ \sqrt{-c_1} \left(\text{Bessel}J(0, 2 \sqrt{-c_1} \sqrt{e^x}) \right)^2 - (h(x, \eta))^2 = 0,$$

(at the conditions $A = 1$ and $-C2 = 0, -C4 = 0, -C1 = 1$).

The solution of the equation (16) which corresponds the function $h(x, \eta)$ is determined from the relation

$$y - h(x, f(x, y, \tau) - x\tau) = 0$$

and can be very complicated.

References:

1. V. Dryuma, *On nonlinear equations associated with developable, ruled and minimal surfaces.* ArXiv:1002.0952 v1 [physycs.gen-ph] 4 Feb.2010 p. 1-13.
2. V. Dryuma, *The Riemann and Einstein-Weyl geometries in theory of differential equations, their applications and all that.* A.B.Shabat et all.(eds.), New Trends in Integrability and Partial Solvability, Kluwer Academic Publishers, Printed in the Netherlands , 2004, p.115–156.